

600360
FAA ADS - 5

FAA ADS - 5



58-P-#1.50

TECHNICAL REPORT


ADS - 5

OZONE MEASUREMENT SURVEY
IN COMMERCIAL JET AIRCRAFT

by

Robert I. Brabets

IIT Research Institute
Technology Center, Chicago 16, Illinois
Under Contract A&DS-608



FEDERAL AVIATION AGENCY

Washington, D.C.

NOVEMBER 1963

TABLE OF CONTENTS

	Page
Summary	1
I. Introduction	3
II. Measuring Techniques	5
III. Results and Discussion	9
A. Source of Ozone	9
B. Differences among Aircraft	9
C. Ozone Measurements	12
D. Internal and External Ozone Concentrations	15
IV. Conclusions	16
Acknowledgments	17
References	19
Appendix - Figures and Tables	21

LIST OF FIGURES

FIGURES		Page
1	Two-Stage Refrigeration System	22
2	Parallel Flights Made on April 23, 1963	23
3	Parallel Flights Made on April 28, 1963	24
4	Record of Ozone Detection (Domestic)	25
5	Record of Ozone Detection (Northern Routes)	26
6	Ozone Detected during Flights Over Middle Latitudes	27
7	Ozone Detected during Flights Over Northern Latitudes	27
8	Profile of Flight of Feb. 14, 1963, and Tropopause Altitude	28
9	Ozone Detection and Tropopause Altitude	29

LIST OF TABLES

TABLE		Page
1	Nominal Comparison of Aircraft Pressurization Systems	30
2	Ozone Detected on Two Different Aircraft during Parallel Flights between Seattle and Chicago on April 9, 1963	31
3	Summary of Data on Ozone Detection during Domestic Flights (Middle Latitudes)	32
4	Summary of Data on Ozone Detection during Flights in Northern Latitudes	33
5	Summary of Data on Ozone Detection during Domestic Flights (Middle Latitudes) in Spring 1962	34
6	Chronological Summary of Ozone Detection during Flights over Middle Latitudes	35
7	Chronological Summary of Ozone Detection during Flights over Northern Latitudes	47
8	Chronological Summary of Ozone Detection during Flights over Polar Latitudes	52
9	Chronological Summary of Ozone Detection Data Obtained on Contract No. FA 2688	53

FEDERAL AVIATION AGENCY
TECHNICAL REPORT ADS-5
OZONE MEASUREMENT SURVEY IN COMMERCIAL JET AIRCRAFT
by Robert I. Brabets

SUMMARY

The purpose of this survey was (1) to accurately measure ozone concentration in commercial jet aircraft cabins and/or flight crew compartments on flights above 25,000 feet, in order to obtain a 12-month statistical evaluation, with emphasis on seasonal and meteorological correlations; and (2) to locate and chart the ozone-enriched air masses in order to obtain further meteorological correlations and to establish any abnormal conditions which result in exposure to large ozone concentrations.

Between September 1, 1962, and August 31, 1963, the ozone concentration in aircraft was measured during 295 commercial jet flights. These flights ranged over all segments of the United States and also included sections of Canada and the North Atlantic. All types of commercial jet aircraft currently employed by air carriers were monitored.

The ozone measurements recorded on each flight were evaluated. All the flights are tabulated in chronological order in the Appendix. The report includes summary tables of the data for the middle and northern latitudes. The data are also correlated graphically to show seasonal variations. The

IIT RESEARCH INSTITUTE

maximum continuous ozone exposure encountered on a domestic flight was 20 or more parts per hundred million by volume for 140 minutes; the maximum on a northern flight was between 20 and 30 PPHM for four hours; the highest concentration encountered was 35 to 40 PPHM for 20 minutes. The most significant finding was that little or no ozone was detected on flights made below the tropopause. At or above the tropopause the internal concentration was usually above 5 PPHM, and in most cases above 10 PPHM.

OZONE MEASUREMENT SURVEY IN COMMERCIAL JET AIRCRAFT

I. INTRODUCTION

Shortly after commercial jet aircraft were placed in operation on regular flight schedules, many passengers reported that the odor of ozone was detectable in the fuselage from time to time. It was doubtful that the ozone was being generated inside the aircraft. However, since air entering the aircraft was required to pass through a compression cycle which raised the temperature to 250°F, ozone, being thermally unstable, could be expected to decompose at this temperature. Equally curious was the fact that ozone was encountered at altitudes of 30,000 to 40,000 feet, whereas the ozone layer of the atmosphere was generally considered to exist at 80,000 feet.

The toxicity and the physiological effects of ozone at certain concentration levels have been reported in the literature, and the reactivity of ozone with certain polymeric materials is also documented.

In December 1961, under the sponsorship of the Federal Aviation Agency, a 4-month study (Contract No. FA2688) was initiated at IIT Research Institute. The purpose was to determine the frequency and the concentration of ozone in commercial jet aircraft.

This study was made on 38 separate flights on the commercial routes of Airline "A". It covered the northern and central portions of the United States from New York to California and included five flights between California and Hawaii. The flights lasted from 60 to 325 minutes and covered altitudes from 9,000 to 39,000 feet. Ozone ranging in concentration from 1 to 22 parts per hundred million (PPHM) was detected during 50% of the total flight time.

On the basis of these preliminary results, the Federal Aviation Agency continued the study for a one-year period (Contract No. ARDS-608) and broadened the scope to include as many types of aircraft and flight patterns as possible. Seven major airlines cooperated by providing free air transportation for the research personnel who were responsible for the ozone measurements. The coast-to-coast routes were covered by flights operated by Airlines "C", "B", and "A"; the southern portion of the United States was covered by Airlines "D" and "C"; the northern latitudes were covered by Airline "E" polar route to London, Airline "F" route to Alaska, and Airline "G" route to Copenhagen. All types of U.S. type certificated jet transport category aircraft were used. The ultimate objective was to correlate the frequency and the concentration of ozone with geographic locations, meteorological conditions, flight patterns, altitudes, and types of aircraft.

II. MEASURING TECHNIQUES

The ozone analyzers used were selected from several commercially available types. The analyzer is based on the microcoulomb ozone sensor conceived by Dr. A. W. Brewer of Oxford University. The basic principle of this sensor is the well-known oxidation-reduction of potassium iodide, which is contained in the sensing solution. Two reactions take place in the sensor: oxidation of potassium iodide by ozone, and reduction of the released iodine by hydrogen. These electrochemical reactions take place at polarized electrodes which are immersed in the sensing solution. The measurement of electron flow or current through the external circuit is directly proportional to the mass of ozone entering the sensor per unit time. A constant-volume solution pump, a constant-volume sample pump, and a calibrated, electronic read-out system complete the instrument package.

The first analyzer used in this study was obtained in 1960 and was calibrated regularly in our laboratories by comparison with the standard wet analytical method in which sodium thiosulfate is used to titrate a buffered potassium iodide solution. This analyzer was also found to agree with instruments of other manufacturers. However, because it requires a 110-volt, a.c. power supply, certain modifications and auxiliary equipment were necessary before it could be operated in an aircraft.

Two additional analyzers were purchased when the scope of the program was broadened. These instruments are powered by rechargeable batteries, are more compact and hence less conspicuous, and are considerably more convenient to handle. The calibration of these instruments was regularly checked at 25-hour intervals against laboratory established ozone concentrations of 2, 8, 16, and 40 PPHM. Additional checks were made when unusual data were recorded. If the instruments did not agree to within 5% of the laboratory value, the previous data were discarded. The calibration was also checked for operation at the subatmospheric pressures found in jets. However, because of aircraft vibrations, slight temperature and pressure fluctuations, and the instruments' response time, the concentrations recorded during flights can be considered accurate only to within 10%.

At the start of the program it was considered advisable to attract the least possible attention from passengers. Consequently, the analyzer was generally placed in the cockpit and started in operation before the passengers boarded the aircraft. The engineer monitoring the analyzer usually rode the observer's seat in the cockpit. His duties were primarily to record pertinent flight data such as altitude and flight path and periodically check the operation of the analyzer.

Since the analyzer was operated in the cockpit, it was necessary to know whether the ozone concentration in the cockpit was the same as that in the main cabin. The air is changed every 3 minutes in the cabin, and every 1-1/2 minutes in the cockpit. It was also necessary to know whether the ozone content of the fresh incoming air differed from that of the air about to be exhausted and whether the air circulation in the aircraft was sufficient to eliminate any stagnant areas.

It was imperative that this information be gained early in the program. Because of cockpit space limitations, it was desirable to monitor some flights with the analyzer positioned in the cabin. Any relationship between ozone content and instrument location had to be determined before the data were evaluated.

Several flights with two or more analyzers aboard the same aircraft were necessary to answer these questions. The ozone content of the air was found to fluctuate rapidly; the concentration could double, or halve, in a matter of several minutes. This knowledge precluded the use of one instrument which could be moved to different locations in the aircraft. The incoming air could be analyzed by an instrument placed a few inches away from the auxiliary ventilating system, or "eyeball." The general level of ozone and the exhaust air could be analyzed by an instrument placed on the flight deck of the aircraft. Further checks could be made by turning the eyeball on and off.

This technique was very successful in establishing standards in the Type "M" and "Q" jets. Both these aircraft have eyeballs in the flight deck located adjacent to a convenient instrument position. The Type "O" jets do not have as convenient a testing location; but since these jets have a faster air turnover rate, it was believed that the results would be comparable.

Comparison of the ozone content of the flight deck with that of the main cabin was accomplished simply by using two analyzers. The second instrument was placed in the cabin before the passengers boarded the aircraft. Not too many tests were required to establish the fact that the ozone content was generally consistent throughout the aircraft. The slower air turnover in the cabin tended to dampen fluctuations, but the average values were the same as those measured in the flight deck. The incoming air, of course, showed the widest fluctuations; but its average value was, for all practical purposes, the same as the others. No areas of stagnation were located, even though sections of the after-cabin occasionally showed a temperature differential of 5°F or more. Smoke, particularly from cigars or pipes, could temporarily reduce the ozone content immediately adjacent to the smoker by as much as 10%. It was also noted that the ozone level adjacent to the galley could drop about 10% during periods when food was being prepared. Even though these variations in ozone content could be considered minor and within normal fluctuations, the

majority of the measurements were made in the cockpit, with the instrument located close to an incoming airstream.

III. RESULTS AND DISCUSSION

A. Source of Ozone

The source of ozone is the external air. This fact is readily apparent from examination of the recordings. Operation of the analyzer for periods of up to one hour before takeoff but with all aircraft equipment operating established the base or zero level for each flight. During many flights, no ozone was detected for various periods of time. As a general rule, no ozone was detected below 25,000 feet, although a few exceptions were noted. For example, ozone was recorded at an altitude of several thousand feet over the Los Angeles smog basin and in areas adjacent to thunderstorms.

B. Differences among Aircraft

It was not possible to determine which type of aircraft would contain more ozone when traveling through a high external concentration. This is apparent from a study of the pressurization system and from data obtained on parallel flights.

The design characteristics of various aircraft pressurization systems was investigated. Specifications for each type of aircraft vary. In addition, further modifications are encountered in the same basic type of aircraft according to the preferences of a particular airline. Table 1 shows that the

pressurization systems differ with respect to compression temperatures and time periods at these temperatures. At first glance it appears that correlation of these factors with internal ozone concentration would give an ozone decomposing factor for each type of aircraft. This factor, in turn, could be used to calculate the ozone concentration of the outside air. Unfortunately, the actual operation of these various pressurization systems precludes any correlation.

When an aircraft cruises at high altitudes, some of the heat of compression is removed from the incoming air to control the internal cabin temperature. All the aircraft have two-stage refrigeration systems similar to the schematic illustration in Figure 1. In addition to the heat removed by the refrigeration system, considerable heat is also removed by conduction and radiation from the ductwork of the ventilating system. During the majority of flights, the primary cooler is regulated to control the cabin temperature. This is accomplished simply and economically by cooling only a portion of the incoming air. The bypassed hot air is mixed with the cooled air to produce the desired temperature in the cabin. This control is easily automated and rapidly corrects for fluctuations in the temperature of the external air, which often varies by 20°F in a matter of minutes.

The secondary refrigeration unit is not required unless the passenger loading is high, the outside temperature is higher than normal, or the flight personnel prefer a cooler cabin. Daytime flights, because of the additional heat supplied by the sun, require more cooling than nighttime flights. Other variables are changes in the outside temperature with altitude and changes in the barometric pressure during the course of a flight. One last feature that eliminates any possibility that internal ozone concentration can be correlated with the pressurization systems is the fact that the aircraft are operated by personnel who have individual preferences. Some crews prefer a warm cabin, some a cool cabin. Some maintain a smaller differential between the inside and outside pressure than others. Some crews anticipate altitude changes and maintain a constant pressure throughout a flight. Sometimes the cabin is cooled a little more during meal times. These differences preclude any calculation of the time that the incoming air is retained in the hot zone during an entire flight.

Numerous attempts were made to determine the relative effectiveness of the various pressurization systems in decomposing ozone. For example, the data listed in Table 2 were obtained on parallel flights between Seattle and Chicago on April 9, 1963. An "N" jet made the flight at 37,000 ft., followed 45 minutes later by a "Q" jet at 33,000 and 37,000 ft.

The pressurization systems of both aircraft were operating in a normal manner, i.e., with part of the incoming air bypassing the primary cooler. Evaluation of these data and comparisons between the data obtained on similar parallel flights (Figures 2 and 3) led to the conclusion that there is no significant difference in ozone-decomposing efficiency among the various types of aircraft.

C. Ozone Measurements

During the period from September 1, 1962, through August 31, 1963, the ozone concentration was monitored on 285 flights, which ranged from short (e.g., Chicago to New York) to medium (e.g., Chicago to San Francisco) to long (e.g., transcontinental, London to San Francisco). The average domestic flight lasted 3-1/4 hours; transcontinental flights over the northern latitudes averaged 7-1/2 hours. The flights were chosen at random, the only controls being to have at least two flights per week and to cover as large an area as possible. During the spring months the number of flights was increased to 6 or 8 per week.

The record for a typical domestic flight is shown in Figure 4. This trace shows that a substantial ozone concentration was encountered shortly after the aircraft reached cruising altitude. As the flight continued westward, the ozone concentration increased and then quite suddenly dropped to zero. No ozone was encountered over the plains states. Then another region of appreciable ozone was encountered. Fifteen minutes before the aircraft started to

descend, the ozone level again dropped to zero. From takeoff to landing, this particular flight lasted 5 hours and 22 minutes, 4 hours and 41 minutes of which were spent at the cruising altitude of 35,000 feet. Ozone in excess of 3 PPHM was encountered for 2-1/2 hours, or 46% of the total flight time. Ozone in excess of 5 PPHM was encountered for 2-1/4 hours, or 42% of the total flight time.

All the flights were evaluated in this manner. An example of a northern flight is presented in Figure 5. The data, assembled in monthly groups, are summarized in Tables 3, 4, and 5. A chronological listing is given in the Appendix. Also included in the Appendix are the data obtained from December 19, 1961, through April 30, 1962, on Contract No. FA 2688.

Examination of the individual flights shows that no ozone was recorded during some of them, whereas appreciable quantities were detected during others. With respect to domestic flights, the highest concentration of ozone detected for the longest period of time occurred in April 1963 on a flight from Seattle to Chicago, when the concentration exceeded 20 PPHM for the entire 140-minute period that the plane was at cruising altitude, 37,000 ft. With respect to the northern latitudes, the highest concentration detected for the longest period also occurred in April 1963, on a flight from London to Seattle, when the concentration fluctuated between 20 and 30 PPHM for 4 hours. The highest

concentration detected during any of the flights occurred in March 1963 on a flight from Anchorage to New York, when a concentration of 35 to 40 PPHM was recorded for a 20-minute period. The concentration was above 30 PPHM for 1 hour before the maximum concentration was recorded.

The expected seasonal variations are quite evident when the data are presented graphically, as in Figures 6 and 7. How the ozone level detected over the middle latitudes differs from that detected over the northern latitudes is clearly shown in the figures.

The ozone data can also be compared with the altitude data. As expected, the frequency and the concentration of ozone generally increases as the flight altitude increases. However, rather than the absolute altitude, the altitude of the tropopause appears to be the deciding factor. Figure 8 is a profile of meteorological conditions which existed during the flight illustrated in Figure 4. When the two figures are combined in Figure 9, it is obvious that ozone is encountered above the tropopause. Very seldom is ozone detectable below the tropopause. Further correlations of the quantity of ozone with height above the tropopause, with variations in latitude, and with seasonal variations are lengthy and complicated. The Meteorology Department of Airline "A" is studying the data for significant trends.

D. Internal and External Ozone Concentrations

No correlation has yet been established between internal and external ozone concentrations. During the latter months of the program, attempts were made to determine whether a constant relationship exists between the ozone concentration inside and outside the aircraft. This problem proved to be more difficult than expected. The data obtained to date are too sparse to provide an adequate correlation.

An approximation is available through comparison with data obtained by the Air Force on the Ozonesonde program. This comparison indicates that the aircraft pressurization system efficiently destroys low concentrations of ozone, but the efficiency decreases as the concentration of ozone increases. In other words, if the external air contains 5 to 10 PPHM ozone, little or none of this ozone will reach the cabin. An external ozone concentration of 50 PPHM will introduce perhaps 10 to 15 PPHM into the cabin, and an external concentration of 75 PPHM might be sufficient to produce the 30 PPHM occasionally recorded in the cabin. This correlation is very tentative and is based on readings from two different types of analyzers. The correlation should be narrowed by making controlled flights in the same area and at the same time as the Ozonesonde balloon ascents, with instrumentation that measures both the internal and the external ozone concentration.

IV. CONCLUSIONS

The principal objectives of the program were attained. The average and the maximum exposures of flight personnel and passengers to ozone were determined.

The ozone concentration varies with altitude, latitude, and season. The ozone-enriched air masses were located and identified as segments of the lower stratosphere. The altitude of the tropopause thus determines whether ozone will be detectable internally. The internal ozone concentration can be considered negligible (1 to 3 PPHM) on flights made below the tropopause. Fluctuations in the altitude of the tropopause are directly correlated with latitude and season.

Thus, only one factor need be determined in order to predict whether a significant concentration of ozone (5 PPHM or more) will be encountered on any projected flight. This one-variable correlation also allows extrapolation of the averaged data to specific routes. For example, those routes which are consistently flown above the tropopause, i.e., polar flights, will have a higher than average ozone exposure. Individual airlines can compare their flight patterns with tropopause data and obtain a better indication of the concentration of ozone likely to be encountered by their personnel. Unfortunately, time was not available to establish the relationship between ozone concentration and distance above the tropopause or seasonal variations in the height of the tropopause.

ACKNOWLEDGMENTS

This is Report No. IITRI-C232-13 (Final Report) of IITRI Project C232, entitled "Ozone Measurement Survey." The report covers the period from August 7, 1962, through September 16, 1963. The work was carried out for the Federal Aviation Agency under Contract No. ARDS-608. Messrs. T. W. Sanford, Jr., and L. S. Jaffe of the FAA served as Project Manager and Alternate Project Manager, respectively.

The invaluable assistance given to IITRI by the personnel of the seven cooperating air lines is gratefully acknowledged. These personnel include:

Mr. S. G. Folsom, Assistant to Division Chief Pilot, Pan American World Airways, New York.

Mr. D. C. Garrett, Jr., Assistant to Executive Vice President, Operations, Delta Air Lines, Inc., Atlanta, Georgia.

Mr. Henry T. Harrison, Director of Meteorology, United Air Lines, Chicago, Illinois.

Mr. C. Hughes, Manager, Flight Administration, Northwest Orient Airlines, St. Paul, Minnesota.

Dr. F. Jonsson, Medical Director, Scandinavian Airlines System, Inc., Stockholm, Sweden.

Dr. G. J. Kidera, Medical Director, United Air Lines, Chicago, Illinois.

Mr. M. I. Lawson, Meteorology Department, United Air Lines, Chicago, Illinois

Dr. L. G. Lederer, Medical Director, American Airlines, New York.

Mr. S. Miller, Division Chief Pilot, Pan American World Airways, New York.

Mr. R. B. Mueller, Staff Vice President, Flight Operations, Trans-World Airlines, Inc., Kansas City, Missouri

IIT RESEARCH INSTITUTE

Mr. T. Nilert, President, Scandinavian Airlines System, Inc.,
New York.

Mr. J. Otto, Operations Manager, Pan American World Airways,
Chicago, Illinois

Dr. O. B. Schrueder, Medical Director, Pan American World
Airways, New York.

Mr. H. Skjoldager, Operations Manager, Scandinavian Airlines
System, Copenhagen, Denmark.

The cooperation of the Mast Development Company of
Davenport, Iowa, in the selection of the ozone analyzer is
gratefully acknowledged.

The following personnel at IITRI assisted in obtaining
the ozone data: Mr. C. K. Hersh, Dr. M. J. Klein, Mr. L. C.
Kinney, Dr. S. Katz, Miss E. J. Grimm, Mr. T. A. Burgwald,
Mr. T. M. Rymarz, Dr. W. M. Whaley, Mr. A. J. Tulis,
Mr. A. J. Gaynor, Dr. I. J. Solomon, Mr. T. A. Erikson,
Mr. J. L. Austing, Mr. E. R. Nielson, and Mr. W. A. Faucette.

REFERENCES

1. Clamman, H. G., Physical and Medical Aspects of Ozone In Benson, O. O. and Strughold, H. (Eds.): Physics and Medicine of the Atmosphere and Space, Chapter IX.
2. Clamann, H. G. and Bancroft, R. W.: Toxicity of Ozone in High Altitude Flight. In Ozone Chemistry and Technology, p. 352, Advances in Chemistry Series, No. 21. American Chemical Society, 1959.
3. Ernsting, J.: Physiologic Aspects of the Significance of Ozone in Pressurized Cabins. I.A.M. Memo No. S-20. RAF Inst. of Av. Med., Farnborough, Sept. 1959.
4. Griswold, S. S., Chambers, L. A., and Motley, H. L.: Report of a Case of Exposure to High Ozone Concentrations for Two Hours. A.M.A. Arch. Ind. Health, 15:108, 1957.
5. Int. J. Air and Water Poll.: The Interaction of Ozone with Plastic and Metallic Materials, 1961, Vol. 4, p. 70-78.
6. King, Maurice, E.: Toxicity of Ozone. V. Factors Affecting Acute Toxicity. Ind. Med. Surg., 32:93, 1963.
7. Kleinfeld, M. and Giel, C. P.: Clinical Manifestations of Ozone Poisoning; Report of a New Source of Exposure. Amer. J. Med. Sci., 231:638, 1956.
8. Lagerwerff, J. M., Kane, G. L., and Thornberg, G. H.: The Effects of Repeated and Prolonged Exposure to High Concentration of Ozone on the Vision of Airline Pilots. Univ. of Minn. Inst. of Technology, Rosemount Aeronautical Laboratories, Res. Report No. 180, May, 1961 (Federal Aviation Agency Contract FA-824).
9. Mittler, S.: Toxicity of Ozone. IV: Silicone Aerosols and Alcohol Vapor Therapy in Ozone Poisoning. Ind. Med. Surg., 27-43, 1958.
10. Mittler, S., Hedrick, D., King, M., and Gaynor, A.: Toxicity of Ozone. I: Acute Toxicity. Ind. Med. Surg., 25: 301, 1956.
11. Mittler, S., Hedrick, D., and Phillips, L.: Toxicity of Ozone. II: Effect of Oxygen and Carbon Dioxide upon Acute Toxicity. Ind. Med. Surg., 26-63, 1957.

12. Mittler, S., King, M., and Burkhardt, B.: Toxicity of Ozone: III. Chronic Toxicity. A.M.A. Arch. Ind. Health, 15:191, 1957.
13. Scheel, L. D., Dobrogorski, O. J., Mountain, J. T., Svirbely, J. L., and Stokinger, H. E.: Physiological, Biochemical, Immunological and Pathological Changes Following Ozone Exposure. J. Appl. Physiol., 14:67, 1959.
14. Stokinger, H. E.: Evaluation of the Hazards of Ozone and Oxides of Nitrogen-Factors Modifying Toxicity. A.M.A. Arch. Ind. Health, 15:181, 1957.
15. Stokinger, H. E.: Factors Modifying Toxicity of Ozone. In Ozone Chemistry and Technology, p. 360, Advances in Chemistry Series, No. 21. American Chemical Society, 1959.
16. Stokinger, H. E., Wagner, W. D., and Dobrogorski, O. J.: Ozone Toxicity Studies. III: Chronic Injury to Lungs of Animals Following Exposure at Low Level. A.M.A. Arch. Ind. Health, 16:514, 1957.
17. Thorp, C. E.: Bibliography of Ozone Technology. Vols. I and II. Chicago, Ill., Armour Research Foundation, J. S. Swift Co. 1954, 1955.

APPENDIX - FIGURES AND TABLES

IIT RESEARCH INSTITUTE




Figure 1
TWO-STAGE REFRIGERATION SYSTEM





Figure 2

PARALLEL FLIGHTS MADE ON April 23, 1963



The numbers encircled are PPHM at the indicated Time over the check point

Figure 3
PARALLEL FLIGHTS MADE ON APRIL 28, 1963




Figure 4

RECORD OF OZONE DETECTION
(New York to San Francisco, February 14, 1963)

IIT RESEARCH INSTITUTE




Figure 5

RECORD OF OZONE DETECTION
(NORTHERN ROUTES)

IIT RESEARCH INSTITUTE




Figure 6

OZONE DETECTED DURING
FLIGHTS OVER MIDDLE LATITUDES





Figure 7

OZONE DETECTED DURING
FLIGHTS OVER NORTHERN LATITUDES



Altitude, ft.

Figure 8

PROFILE OF FLIGHT OF FEBRUARY 14, 1963,
AND TROPOPAUSE ALTITUDE




Figure 9

OZONE DETECTION AND TROPOPAUSE ALTITUDE

Table 1
NOMINAL COMPARISON OF AIRCRAFT PRESSURIZATION SYSTEMS

	Aircraft "M"	Aircraft "N"	Aircraft "O"	Aircraft "P"	Aircraft "Q"
Mode of pressurization: Normal Alternate	Turbocompressor Bleed air, 7th stage	Turbocompressor Bleed air, 7th stage	Turbocompressor Bleed air, 17th stage	Turbocompressor Bleed air, 17th stage	Turbocompressor None
Location of air inlet	Inboard engine pods	Inboard engine pods	Bottom of fuselage, forward of wings	Bottom of fuselage, forward of wings	Lower front of fuselage
Fuselage volume	9400 cu ft	8400 cu ft	4500 cu ft	5300 cu ft	9300 cu ft
Avg. time per air change: In flight deck In cabin	1.5 min 3 min	1.5 min 3 min	1 min 2 min	1.25 min 2.5 min	1.5 min 3 min
Volume of ductwork upstream of prime cooler	12 cu ft	12 cu ft	0.25 cu ft	0.25 cu ft	1.1 cu ft
Ductwork material	Type 321 SS	Type 321 SS	SS	SS	Aluminum
Airflow per turbo- compressor at 35,000 ft	1600 cfm	1600 cfm	830 cfm	830 cfm	700 cfm
Pressure in duct at 35,000 ft	18 psia	18 psia	14 psia	14 psia	14 psia
Temperature in duct at 35,000 ft	250-275°F	250-275°F	290°F	290°F	200°F
Residence time of air in hot zone	0.45 sec	0.45 sec	0.02 sec	0.02 sec	0.1 sec

MIT RESEARCH INSTITUTE

Table 2
OZONE DETECTED ON TWO DIFFERENT AIRCRAFT
DURING PARALLEL FLIGHTS BETWEEN SEATTLE AND CHICAGO
ON APRIL 9, 1963

Check Point	Aircraft "N"		Aircraft "Q"	
	Altitude, ft	Ozone Conc., PPHM	Altitude, ft	Ozone Conc., PPHM
Spokane	37,000	14	33,000	5
Mullan Pass	37,000	8	33,000	8
Billings	37,000	10	33,000	10
Dupre	37,000	12	33,000	5
Sioux Falls	37,000	4	37,000	3
Mason City	37,000	5	37,000	3
Dubuque	37,000	5	37,000	5

--- RESEARCH INSTITUTE

Table 3
SUMMARY OF DATA ON OZONE DETECTION
DURING DOMESTIC FLIGHTS (MIDDLE LATITUDES)

Month	Number of Flights	Total Flight Time, min	Time above 25,000 ft, min	Time Ozone Detected, min									
				>1 PPHM	>3 PPHM	>5 PPHM	>10 PPHM	>15 PPHM	>20 PPHM	>25 PPHM	>30 PPHM		
Sept 62	11	2269	1690* (74.5)	513 (22.6)	153 (6.7)	73 (3.0)							
Oct 62	11	2257	1731 (76.7)	265 (11.7)	80 (3.6)	10 (0.4)							
Nov 62	17	3938	3002 (76.2)	1232 (31.4)	290 (7.4)	32 (0.8)							
Dec 62	13	2427	1787 (73.6)	745 (30.6)	235 (9.7)	150 (6.2)	25 (1.0)						
Jan 63	25	4566	3357 (73.5)	748 (16.4)	443 (9.7)	730 (8.1)	110 (2.4)	85 (1.8)	70 (1.5)	50 (1.1)	10 (0.2)		
Feb 63	18	3168	2237 (70.6)	1048 (33.1)	810 (17.7)	610 (13.3)	250 (5.5)						
Mar 63	30	5840	4419 (75.6)	2569 (44.0)	1903 (32.5)	1554 (26.6)	785 (13.4)	275 (4.7)	35 (0.6)				
Apr 63	26	4758	3451 (72.5)	2396 (50.3)	1927 (40.5)	1715 (36.1)	1125 (23.7)	904 (19.0)	644 (13.5)	290 (6.1)	105 (2.2)		
May 63	24	4689	3391* (72.3)	1712 (36.5)	871 (18.5)	650 (13.8)	198 (4.2)	70 (1.5)	15 (0.3)				
June 63	23	3660	2476 (67.0)	1194 (32.6)	370 (10.0)	287 (7.8)	110 (3.0)	40 (1.1)					
July 63	11	1655	1134 (68.4)	59 (35.9)	71 (4.3)	56 (3.4)	21 (1.2)						
Aug 63	19	3035	1897 (62.5)	610 (20.1)	30 (1.0)	5 (0.2)							

* Numbers in parentheses refer to percent of total flight time.

IIT RESEARCH INSTITUTE

Table 4
SUMMARY OF DATA ON OZONE DETECTION
DURING FLIGHTS IN NORTHERN LATITUDES

Month	Number of Flights	Total Flight Time, min	Time above 25,000 ft, min	Time Ozone Detected, min							
				>1 PPHM	>3 PPHM	>5 PPHM	>10 PPHM	>15 PPHM	>20 PPHM	>25 PPHM	>30 PPHM
Jan 63	8	3628	3105* (85.5)	2483 (63.4)	1904 (52.5)	1548 (47.7)	937 (25.8)	285 (7.8)	100 (2.7)	30 (0.8)	15 (0.4)
Feb 63	5	2544	2253 (88.6)	1761 (69.3)	1461 (57.5)	1131 (44.4)	601 (23.6)	260 (10.2)	85 (3.3)		
Mar 63	11	4666	4104 (88.1)	3657 (79.2)	3311 (71.2)	3151 (67.7)	2273 (48.8)	1433 (30.7)	744 (16.0)	454 (9.7)	175 (3.7)
Apr 63	6	3166	2833 (89.3)	2753 (87.0)	2341 (73.8)	2110 (66.7)	1804 (57.0)	1276 (40.3)	716 (22.5)	191 (6.0)	86 (2.7)
May 63	9	3441	2962 (86.2)	2632 (76.5)	1524 (44.3)	1419 (41.3)	1109 (32.3)	839 (24.4)	587 (17.0)	215 (6.2)	30 (0.9)
June 63	7	3374	2960 (87.7)	2105 (62.3)	1835 (54.3)	1735 (51.3)	1148 (34.0)	624 (18.5)	152 (4.5)		
July 63	4	1330	1113 (83.3)	796 (59.8)	604 (45.5)	589 (44.3)	481 (36.1)	302 (22.7)	95 (7.1)		
Aug 63	4	1271	1089 (85.3)	595 (46.8)	320 (25.2)	265 (20.8)	45 (3.5)				

* Numbers in parentheses refer to percent of total flight time.

Table 5
SUMMARY OF DATA ON OZONE DETECTION
DURING DOMESTIC FLIGHTS (MIDDLE LATITUDES) IN SPRING 1962

Month	Number of Flights	Total Flight Time, min	Time above 25,000 ft, min	Time Ozone Detected, min				
				> 1 PPHM	> 3 PPHM	> 5 PPHM	> 10 PPHM	> 15 PPHM
Feb	14	2880	2375 (82.5)*	930 (32.2)	380 (13.2)	270 (9.4)	185 (6.4)	5 (0.2)
Mar	14	3095	2495 (80.7)	1885 (60.8)	595 (19.2)	390 (12.6)	115 (3.7)	
Apr	8	790	435 (55.1)	425 (53.8)	150 (19.0)	125 (15.8)	115 (15.1)	85 (10.7)
								10 (1.2)

* Numbers in parentheses refer to percent of total flight time.

Table 6
CHRONOLOGICAL SUMMARY OF OZONE DETECTION DURING FLIGHTS OVER MIDDLE LATITUDES

Date	Flight Route		Time of Day	Flight Time, min	Altitude, ft	Time at Altitude, min	Time Ozone Detected, min							
	Origin	Destination					>1 PPM	>3 PPM	>5 PPM	>10 PPM	>15 PPM	>20 PPM	>25 PPM	>30 PPM
9/17/62	Chicago	S. F.	D	232	28,000 31,000	31 163	5							
9/17/62	S. F.	Honolulu	D	286	28,000 31,000	139 94	15							
9/19/62	Honolulu	S. F.	N	270	33,000	215								
9/18/62	S. F.	Chicago	D	215	29,000 37,000	80 80	25							
9/27/62	Chicago	New York	D	103	37,000 39,500	15 33	15							
9/27/62	New York	S. F.	D	314	31,000 39,000	148 115	70 115	30 30	15 20					
9/23/62	New York	Chicago	D	105	31,000 35,000	18 30	18 30	18 30	13 25					
10/4/62	Chicago	New York	D	105	25,000 27,000	35 25	15							
10/4/62	New York	S. F.	D	308	28,000 31,000 35,000	73 143 45	60 45	15 15						
10/5/62	New York	Denver	D	202	28,000 31,000	7 158	50	15						
10/3/62	Denver	Chicago	D	129	25,000 29,000 31,000	18 10 15								
10/11/62	Chicago	New York	D	94	29,000	44								
10/11/62	New York	L. A.	D	315	28,000	262								
10/11/62	L. A.	Denver	N	100	27,000	63								
10/11/62	Denver	Chicago	N	105	29,000 31,000	34 35								
10/19/62	Chicago	S. F.	D	230	31,000 35,000	80 105	60							
10/19/62	S. F.	Honolulu	D	270	31,000 35,000	180 47								

Table 6 (Cont.)

Date	Flight Route		Time of Day	Flight Time, min	Altitude, ft.	Time at Altitude, min	Time Ozone Detected, min								
	Origin	Destination					> 1 PPM	> 1 PPM	> 5 PPM	> 10 PPM	> 15 PPM	> 20 PPM	> 25 PPM	> 30 PPM	
10/20/62	Honolulu	S. F.	N	273	31,000	225	40								
10/20/62	S. F.	Chicago	D	235	27,000	150									
10/25/62	Chicago	New York	D	92	29,000	47	45	20							
10/25/62	New York	S. F.	D	322	28,000	273	90	60	10						
10/26/62	S. F.	Chicago	D	221	27,000	173	30								
11/1/62	Chicago	New York	D	93	29,000	46	40								
11/1/62	New York	S. F.	D/W	317	35,000	165	165	25							
					39,000	103	25								
11/4/62	Chicago	Baltimore	D	105	29,000	45	35	10							
11/5/62	Baltimore	S. F.	N	313	31,000	264	30	5							
11/7/62	S. F.	Baltimore	D	275	29,000	118	35								
					33,000	103	103	43	25						
11/7/62	Baltimore	L. A.	N	301	31,000	165	120	35							
					35,000	67	67	57	7						
11/16/62	L. A.	Baltimore	D	247	33,000	47	45	15							
					37,000	140	120								
11/16/62	Baltimore	S. F.	N	320	28,000	100	30								
					31,000	140	130	55							
					35,000	22	22								
11/17/62	S. F.	Chicago	D	210	29,000	85	35	15							
					33,000	68	45								
11/20/62	Chicago	New York	D	88	39,000	33	30	20							
11/20/62	New York	S. F.	N	336	28,000	297	65								
11/21/62	S. F.	Chicago	D	207	29,000	35									
					33,000	31	5								
					37,000	87	65	10							
11/28/62	Chicago	New York	D	95	29,000	49									
11/28/62	New York	S. F.	N	335	28,000	257	10								
11/29/62	S. F.	Chicago	D	221	29,000	101	10								
					37,000	66									
11/30/62	Chicago	S. F.	D	225	28,000	30									
					31,000	145									

IIT RESEARCH INSTITUTE

Table 6 (Cont.)

Date	Flight Route		Time of Day	Flight Time, min	Altitude, ft	Time at Altitude, min	Time Opone Detected, min								
	Origin	Destination					>1 PPH	>3 PPH	>5 PPH	>10 PPH	>15 PPH	>20 PPH	>25 PPH	>30 PPH	
11/30/62	S. F.	Honolulu	D	250	31,000 35,000 39,000	13 95 85									
12/3/62	Honolulu	L. A.	D	290	29,000	245	180								
12/3/62	L. A.	Denver	H	130	27,000	120									
12/3/62	Denver	Chicago	H	120	29,000	70	50								
12/6/62	Chicago	Seattle	D	226	28,000	184									
12/7/62	Seattle	Chicago	D	194	33,000	121	80								
12/10/62	Chicago	Seattle	D	245	28,000 31,000	115 88	60								
12/11/62	Seattle	Chicago	D	206	37,000	152	75	50	30						
12/19/62	Chicago	Las Vegas	D	188	31,000	151	5								
12/20/62	Las Vegas	Chicago	H	175	27,000 37,000	15 110	95	15							
12/21/62	Chicago	L. A.	H	212	35,000 39,000	39 121	20 120	20 120	10 95	25					
12/21/62	L. A.	Chicago	D/H	215	29,000	159									
12/27/62	Chicago	New York	D	108	25,000	26									
12/27/62	New York	Chicago	H	118	26,000 35,000	14 57	5 55	30	15						
1/3/63	Chicago	Seattle	H	260	31,000 35,000	25 165									
1/8/63	Chicago	L. A.	D	226	31,000	182									
1/10/63	Chicago	Mem. City	D	227	31,000 39,000	46 135	45	10							
1/10/63	Mem. City	Chicago	H	225	33,000 37,000	10 150	45	5							
1/11/63	L. A.	Chicago	D	192	29,000	133									
1/14/63	Chicago	Mem. City	D	217	25,000 31,000	92 89	75 35								

IIT RESEARCH INSTITUTE

Table 6 (Cont.)

Date	Flight Route		Time of Day	Flight Time, min	Altitude, ft	Time at Altitude, min	Time Onset Detected, min					
	Origin	Destination					> 1 PM	> 1 PM	> 1 PM	> 1 PM	> 1 PM	> 1 PM
1/15/63	Man. City	Chicago	N	180	31,000	20						
					33,000	116	15					
1/17/63	Chicago	S. F.	D	233	24,000	59						
					29,000	10						
					31,000	126	20					
1/17/63	Chicago	S. F.	D	235	37,000	29						
					39,000	143	143	143	110			
1/17/63	S. F.	Denver	D	118	27,000	75						
1/17/63	Denver	Baltimore	N	156	37,000	110	75	65	55			
1/18/63	New York	Chicago	D	124	26,000	35						
					33,000	35						
1/21/63	Chicago	Denver	D	125	31,000	90						
1/21/63	Denver	Chicago	D	120	29,000	53						
1/23/63	L. A.	Chicago	N	167	33,000	47	60	15				
					37,000	95						
1/23/63	Chicago	Seattle	N	238	28,000	190	5					
1/24/63	Chicago	L. A.	N	220	31,000	189						
1/26/63	L. A.	Chicago	D	185	29,000	62						
					37,000	78	75	70	70	10		
1/29/63	Seattle	Chicago	N	180	37,000	140	135	135	135	85	70	50
1/29/63	Chicago	Phoenix	D	191	28,000	55						
					31,000	99						
1/29/63	Phoenix	Chicago	D/N	154	37,000	78						
1/31/63	Chicago	New Or.	D	116	31,000	72						
1/31/63	New Or.	Baltimore	N	112	33,000	67						
1/31/63	Chicago	Seattle	D	236	28,000	201	20					
1/31/63	Seattle	S. F.	D	107	33,000	54						

IIT RESEARCH INSTITUTE

Table 6 (Cont.)

Date	Flight Route		Time of Day	Flight Time, min	Altitude, ft	Time at Altitude, min	Time Ozone Detected, min								
	Origin	Destination					>1 PPM	>3 PPM	>5 PPM	>10 PPM	>15 PPM	>20 PPM	>25 PPM	>30 PPM	
2/1/63	S. F.	Chicago	N	193	33,000	127	40								
2/1/63	Wash.	Chicago	D	103	35,000	46									
2/8/63	Chicago	Boston	D	101	33,000	58	50	20	20						
2/8/63	Boston	Chicago	N	123	35,000	42	35	35	5						
					39,000	30	25	25							
2/10/63	Chicago	San Diego	N	222	28,000	183									
2/12/63	L. A.	Chicago	D	214	29,000	98	5								
					37,000	65	65	65	30						
2/14/63	Chicago	New York	D	88	37,000	42	42	5	5						
2/14/63	New York	S. F.	D	330	31,000	290	130	115	90	65					
2/14/63	New York	S. F.	D	322	35,000	281	195	150	135						
2/15/63	S. F.	Chicago	D	223	37,000	159	150	150	125	95					
2/22/63	Chicago	L. A.	D	220	33,000	42	25	15	15						
					39,000	125	125	95	95	60					
2/22/63	L. A.	Chicago	D/N	192	37,000	135									
2/23/63	Las Vegas	Chicago	N	172	28,000	137									
2/28/63	L. A.	Chicago	D	209	29,000	30									
					33,000	120									
2/28/63	Chicago	New York	D	91	33,000	36	20	15							
2/28/63	Chicago	Hartford	D	98	37,000	38	45	45	30						
2/28/63	New York	Chicago	D	117	28,000	72	15								
2/28/63	Hartford	Chicago	D/N	150	35,000	81	81	75							
3/1/63	Chicago	New OrL.	D	111	26,000	78									
3/2/63	New OrL.	Chicago	N	110	27,000	59									
3/4/63	Chicago	L. A.	D	205	31,000	167	25								
3/4/63	Chicago	L. A.	N	225	39,000	166	135	1	110	95					

IIT RESEARCH INSTITUTE

Table 6 (Cont.)

Date	Flight Route		Time of Day	Flight Time, min.	Altitude, ft.	Time at Altitude, min.	Time Ozone Detected, min.						
	Origin	Destination					>1 PPM	>3 PPM	>5 PPM	>10 PPM	>15 PPM	>20 PPM	>25 PPM
3/4/63	L.A.	New York	N	259	37,000	200	145	50	40	20			
3/5/63	New York	Chicago	D	115	39,000	50	50	35	25	25	10	10	
3/8/63	Chicago	L.A.	N	220	31,000	180							
3/8/63	L.A.	Chicago	D	193	37,000	130	130	130	95	15			
3/9/63	Chicago	Seattle	D	221	35,000	10	10	10	10				
					39,000	163	163	163	133	100	40	10	
3/11/63	Seattle	Chicago	D	201	37,000	57	57	57	57	57	32		
					35,000	93	93	93	93	93	93	10	
3/13/63	L.A.	Detroit	N	272	37,000	219	219	185	155	10			
3/16/63	New York	Chicago	D	128	35,000	50							
					28,000	22							
3/18/63	Chicago	Seattle	D	223	39,000	177	177	155	175	85	45	5	
3/18/63	Chicago	L.A.	D	219	35,000	173	120	90	75	15			
3/19/63	Seattle	Chicago	D	195	33,000	15							
					37,000	130	130	100	90	45	10		
3/19/63	L.A.	Chicago	N	198	37,000	146	146	146	136	15			
3/21/63	Chicago	S. F.	N	223	35,000	34	15	5					
					39,000	145	90	15					
3/22/63	S.F.	New York	D	271	29,000	134							
					37,000	112	55	40	40				
3/22/63	New York	Chicago	N	103	31,000	67	20						
3/22/63	Chicago	S. F.	D	230	31,000	126	10	10					
					35,000	47	45	35					
3/22/63	S. F.	Honolulu	N	285	4,000	256	100	10					
3/24/63	Honolulu	S. F.	D	260	33,000	75							
					37,000	140	45	45	15				
3/25/63	S. F.	Chicago	D	202	37,000	153	80	70	55				
3/26/63	Chicago	Seattle	N	222	39,000	180	180	180	180	180	45		

IIT RESEARCH INSTITUTE

Table 6 (Cont.)

Date	Flight Route		Time of Day	Flight Time, min	Altitude, ft	Time at Altitude, min	Time Ozone Detected, min										
	Origin	Destination					>1 PPM	>3 PPM	>5 PPM	>10 PPM	>15 PPM	>20 PPM	>25 PPM	>30 PPM			
3/26/63	Chicago	New Or.	D	107	31,000	65	25	15									
3/27/63	Seattle	Chicago	D	190	37,000	144	144	144	100	30							
3/28/63	New Or.	Chicago	M	125	29,000	73	-										
3/28/63	Chicago	Portland	N	228	28,000	92	5										
							31,000	85	10	10							
3/30/63	New York	Chicago	N	105	28,000	53	45										
					31,000	16	10										
3/31/63	L. A.	Chicago	M	194	37,000	137	90										
4/1/63	Chicago	L. A.	D/M	230	28,000	177	30	25	20								
4/2/63	L. A.	Chicago	D/M	196	37,000	123	95	65	65	55	35	10					
							39,000	12	12	12							
4/2/63	Chicago	Spokane	D	200	39,000	145	145	145	120	95	85	70	25				
4/3/63	Seattle	Chicago	D	210	37,000	138	105	90	80	70							
4/5/63	Chicago	Seattle	D	232	39,000	170	170	90	15								
4/5/63	Portland	Honolulu	M	338	35,000	138	138	123	108								
							39,000	43	20								
					41,000	111											
4/9/63	Seattle	Chicago	D	179	37,000	134	134	115	100	35	15						
4/9/63	Seattle	Chicago	D	191	33,000	93	93	93	78	10							
							37,000	47	47	15							
4/10/63	Chicago	Miami	D	133	37,000	89	15										
4/10/63	Miami	Chicago	M	155	39,000	113	10										
4/15/63	Chicago	New York	D	95	37,000	18	18	18	13	6	6						
							41,000	27	27	27	27	12					
4/15/63	Chicago	New York	D	93	29,000	53											
4/15/63	New York	Minneapolis	N	144	39,000	93	70	55	55	45	35	5					
4/15/63	Minneapolis	Chicago	M	58	33,000	5											

IIT RESEARCH INSTITUTE

Table 6 (Cont.)

Date	Flight Route		Time of Day	Flight Time, min	Altitude, ft	Time at Altitude, min	Time Ozone Detected, min							
	Origin	Destination					>1 PPM	>2 PPM	>5 PPM	>10 PPM	>15 PPM	>20 PPM	>25 PPM	>30 PPM
4/17/63	Chicago	L. A.	D/W	237	28,000	260	185	85	70	30	20			
4/18/63	L. A.	Chicago	D/W	208	37,000	145	145	135	135	120	110	75	20	
4/19/63	Chicago	Baltimore	D	13	37,000	28	5							
4/19/63	Baltimore	Chicago	W	106	28,000 35,000	10 44	35	20	20					
4/22/63	Chicago	Seattle	D	220	39,000	178	145	145	145	115	105	75	25	
4/22/63	Chicago	Portland	W	222	31,000 35,000	68 117	45 117	35 117	35	117	117	70		
4/23/63	Portland	Chicago	D	193	37,000	136	136	136	136	136	125	110	35	15
4/23/63	Seattle	Chicago	D	196	37,000	134	134	134	134	119	99	89	30	
4/23/63	Seattle	Chicago	D	199	37,000	140	140	140	140	140	140	140	70	70
4/26/63	L. A.	Chicago	W	191	33,000 37,000	64 86	40 30	15 15	15					
4/26/63	Chicago	San Diego	W	241	45,000 39,000	175 17	100	45	45	5				
4/30/63	Chicago	S. F.	W	218	31,000	180	10							
5/1/63	S. F.	Chicago	D	211	37,000	154	100	85	50					
5/1/63	Las Vegas	Chicago	D	171	37,000 39,000	31 87	31 87	31 87	25 87	35				
5/4/63	Chicago	Max. City	D	197	39,000	125	50							
5/4/63	Max. City	Chicago	D	201	33,000 37,000 39,000	51 59 40	15							
5/7/63	Chicago	L. A.	D	220	35,000 39,000	93 89								
5/8/63	L. A.	Chicago	D/W	209	37,000	144	5							

IIT RESEARCH INSTITUTE

Table 6 (Cont.)

Date	Flight Route		Time of Day	Flight Time, M. S.	Altitude, ft.	Time at Altitude, min.	Time Ozone Detected, min.						
	Origin	Destination					<1 PMH	2-1 PMH	>1 PMH	>10 PMH	>11 PMH	>20 PMH	>23 PMH
5/10/63	Seattle	Chicago	D	200	33,000	129							
5/10/63	Chicago	Max. City	D	199	31,000	156							
5/11/63	Max. City	Chicago	D	193	37,000	146	5						
5/13/63	Chicago	S. F.	D/M	234	28,000	131							
					31,000	23							
					33,000	30	30	15	15				
5/14/63	S. F.	New York	D	277	33,000	69	65	60	40	5			
					37,000	155	65	60	55	35			
5/14/63	New York	Chicago	D	109	28,000	70	10						
5/15/63	Chicago	Seattle	D	220	39,000	173	160	125	115	75	50	15	
5/19/63	Chicago	Baltimore	D	75	37,000	11	18	18	18	12			
					41,000	18							
5/21/63	Chicago	Baltimore	D/M	75	29,000	33	33						
5/21/63	Baltimore	L. A.	D	289	31,000	245	50	20					
5/22/63	L. A.	Chicago	D	194	37,000	37	60	50	25	15	10		
					39,000	93							
5/22/63	Washington	L. A.	D	293	28,000	255	205	65	35				
5/23/63	Portland	S. F.	D	80	27,000	40	10						
5/23/63	Chicago	L. A.	D	230	35,000	170	170	35	35				
5/24/63	L. A.	Chicago	D	195	33,000	100	90						
					37,000	15	15	10					
3/24/63	L. A.	Chicago	M	192	37,000	126	126	20	20	5			
					33,000	17	17						
5/28/63	Chicago	L. A.	D	219	39,000	161	140	90	65				
5/28/63	L. A.	Chicago	D	206	37,000	135	100	70	40				
6/3/63	Chicago	Baltimore	D	82	33,000	37	30	20	15				
6/3/63	Baltimore	Chicago	D	90	31,000	36	36	5	5				
6/7/63	Chicago	Dallas	D	94	28,000	42							

IIT RESEARCH INSTITUTE

Table 6 (Cont.)

Date	Flight Route		Time of Day	Flight Time, min	Altitude, ft	Time at Altitude, min	Time Ozone Detected, min									
	Origin	Destination					>1 PPM	>3 PPM	>5 PPM	>10 PPM	>15 PPM	>20 PPM	>25 PPM	>30 PPM		
6/7/63	Dallas	Orlando	D	127	24,000 37,000	22 63	22 63	13								
6/9/63	L. A.	Chicago	D/N	189	37,000	136	90	40	30	15						
6/10/63	Tampa	Chicago	D	122	31,000	85										
6/12/63	Chicago	Seattle	D	230	28,000 31,000	40 139	30									
6/12/63	Seattle	New York	N	252	37,000 41,000	135 61	61									
6/13/63	New York	Chicago	D	106	28,000	62	35									
6/13/63	Chicago	Seattle	D	228	28,000	197	65									
6/13/63	Chicago	Baltimore	D	85	33,000	35	15									
6/13/63	Baltimore	Chicago	D	91	31,000 35,000	17 19										
6/15/63	Portland	Chicago	D	195	37,000 39,000	107 44	100 44	44	24							
6/19/63	Chicago	L. A.	D	216	31,000 35,000	48 119	40 85	15 30	30	20						
6/21/63	S. F.	New York	D	280	33,000 37,000	44 185	25 155	15 95	15 95	55	40					
6/21/63	New York	Chicago	D	110	31,000 35,000	27 33	15 33	33	33							
6/21/63	Las Vegas	Chicago	D	166	33,000	122	105									
6/21/63	L. A.	Omaha	D	152	37,000	108	45	20								
6/23/63	Chicago	New York	D	112	37,000 41,000	12 28	12 28	12 28	12 28	20						
6/24/63	Newark	Chicago	D	113	39,000	51										

IIT RESEARCH INSTITUTE

Table 6 (Cont.)

Date	Flight Route		Time of Day	Flight Time, MIN	Altitude, ft	Time at Altitude, MIN	Time Ozone Detected, MIN				
	Origin	Destination					>1 PPM	>3 PPM	>5 PPM	>10 PPM	>25 PPM
6/25/63	Chicago	L. A.	D	219	35,000 39,000	55 106	25				
6/26/63	L. A.	Chicago	D	204	37,000	123	15				
6/28/63	L. A.	Chicago	D/N	197	33,000 37,000 39,000	48 39 51	20				
7/8/63	Pittsburg	Boston	D	62	25,000	29					
7/9/63	Chicago	Denver	D	116	28,000 31,000 35,000	11 24 38					
7/12/63	London	Athens	D	183	33,000	122	80	30	30	5	
7/14/63	Chicago	Portland	D/N	213	26,000	180	15				
7/16/63	Athens	Munich	D	118	29,000	87	10				
7/17/63	Seattle	L. A.	D	135	31,000 37,000	32 51	51	36	26	16	
7/18/63	L. A.	Chicago	N	193	31,000 37,000	11 132	100				
7/21/63	Seattle	L. A.	D	122	31,000	99	85	5			
7/24/63	Las Vegas	Chicago	N/D	183	37,000 41,000	61 48	30				
7/25/63	Chicago	L. A.	D	215	31,000	158	158				
7/26/63	New York	Chicago	D	95	31,000	51					
8/1/63	Chicago	S. F.	N	236	31,000 35,000 37,000	89 82 20	10 75 20				
8/1/63	S. F.	L. A.	D	54	29,000	15					

IIT RESEARCH INSTITUTE

Table 6 (Cont.)

Date	Flight Route		Time of Day	Flight Time, min	Altitude, ft	Time at Altitude, min	Time Ozone Detected, min								
	Origin	Dest. Station					>1 PPM	>3 PPM	>5 PPM	>10 PPM	>15 PPM	>20 PPM	>25 PPM	>30 PPM	
8/1/63	L. A.	Chicago	D	190	33,000	139	85								
8/2/63	Chicago	Miami	D	144	37,000	84									
8/2/63	Miami	Chicago	M	153	35,000	97	45								
8/4/63	Seattle	Chicago	D	177	37,000	134	110	20	5						
8/6/63	Chicago	L. A.	D	214	39,000	148	130	10							
8/12/63	Chicago	Mex. City	D	202	31,000 39,000	112 41									
8/13/63	Mex. City	Chicago	D	201	31,000 37,000	17 137	10								
8/14/63	Chicago	Dallas	M	100	39,000	50	16								
8/14/63	Dallas	New Orl.	M	56	29,000	10									
8/15/63	New Orl.	San Juan	M/D	200	37,000	130									
8/15/63	San Juan	New Orl.	D	200	35,000	138									
8/15/63	New Orl.	Dallas	D	59	28,000	16									
8/15/63	Dallas	Chicago	D	95	28,000 33,000	45 13									
8/19/63	Chicago	L. A.	D	216	28,000	163	30								
8/27/63	L. A.	Chicago	D	196	37,000	130	130								
8/23/63	L. A.	Atlanta	D	227	29,000 33,000	10 177									
8/24/63	Atlanta	Chicago	D	115	26,000 29,000	12 24									

IIT RESEARCH INSTITUTE

Table 7

CHRONOLOGICAL SUMMARY OF OZONE DETECTION DURING FLIGHTS OVER NORTHERN LATITUDES

Date	Flight Route		Time of Day	Flight Time, min	Altitude, ft	Time at Altitude, min	Time Ozone Detected, min									
	Origin	Destination					>1 PPM	>2 PPM	>5 PPM	>10 PPM	>15 PPM	>20 PPM	>25 PPM	>30 PPM		
1/3/63	Seattle	London	D/N	584	27,000 29,000 34,000 38,000 41,000	34 54 63 164 187	10 63 164 187	45	100	30						
1/5/63	London	Chicago	D/N	500	29,000 32,000 35,000 37,000	27 249 82 81	249 82 81	55 20 81	30	81	30					
1/18/63	S. F.	Winnipeg	D	181	29,000	145	100	40	25							
1/18/63	Winnipeg	London	N	447	34,000 39,000	135 262	135 262	135 262	125 262	70 262	125	30	5			
1/23/63	London	L. A.	D	667	32,000 35,000 39,000 40,000	33 285 140 145	155 140 145	130 75 145	115 30 145	115 15 110	55 35					
1/24/63	Seattle	Montreal	D/N	244	33,000	217	185	90	90	50						
1/25/63	Montreal	London	N	434	33,000	280	170	120	120							
1/28/63	London	Seattle	D	571	32,000 35,000 40,000	25 247 250	155 200	155 200	105 200	10 170	70	70	25	15		
2/7/63	Chicago	London	N	427	33,000 37,000	53 304	45 290	45 290	35 245	25 145	95	45				
2/10/63	London	S. F.	D	596	31,000 32,000 35,000 36,000	92 70 96 280	60 70 55 80	30 35 80	30							
2/17/63	Chicago	London	N	453	33,000 34,000 35,000	145 25 225	140 25 225	20 25 195	20 25 80	25 25 80						
2/25/63	Chicago	London	N	425	33,000 34,000 35,000 37,000	31 16 14 367	30 15 230	30 15 215	30 15 140	40	15					

IIT RESEARCH INSTITUTE

Table 7 (Cont.)

Date	Flight Route		Time of Day	Flight Time, min	Altitude, ft	Time at Altitude, min	Time Ozone Detected, min									
	Origin	Destination					>1 PPM	>3 PPM	>5 PPM	>10 PPM	>15 PPM	>20 PPM	>25 PPM	>30 PPM		
2/27/63	London	L. A.	D	643	32,000	20	175	160	150	60						
					35,000	234										
					36,000	161	161	161	161	80	35					
					40,000	180	160	160	135	70	5					
3/10/63	Chicago	London	N	438	33,000	79	60	50	50							
					37,000	302	302	270	270	190	80	10				
3/13/63	London	L. A.	D	7	32,000	24	10									
					35,000	409	409	305	220	135	50	15				
					36,000	182	182	182	182	155	140	70	30			
3/14/63	Chicago	Anchorage	N	367	31,000	47	18	18	15							
					35,000	185	165	165	165	100	85	70	10			
					36,000	69	69	69	69	69	69	39	35			
					39,000	25	25	25	25	25	25	25	25	5		
					40,000											
3/15/63	Anchorage	New York	D/N	377	33,000	95	85	69	69	69	35	30				
					34,000	102	102	42	42	22	22	15				
					37,000	95	95	95	95	85	45	30	20			
					38,000	45	45	45	45	45	45	45	40	40		
3/20/63	Chicago	London	N	448	33,000	89	89	39	39	50						
					37,000	304	235	235	190	160	70	20	10			
3/24/63	London	S. F.	D	658	29,000	54	229	229	215	155	35					
					37,000	229	204	204	204	175	55	15				
					36,000	204	123	123	123	75	45					
					39,000	123										
3/28/63	Seattle	Anchorage	D	160	33,000	128	128	128	120	110	95	30				
3/28/63	Chicago	London	N	435	33,000	173	95	65	65	60	60					
					34,000	18	18	18	18	18	13	13				
					35,000	194	194	155	155	125	100	40	40			
3/29/63	Anchorage	New York	O/N	402	29,000	30	15									
					33,000	72	72	72	72	42	31					
					34,000	186	86	86	86	86	86	35	15			
					38,000	87	87	87	87	87	87	87	72	35		
3/30/63	London	Keflavik	D	137	31,000	55	45	45	45							
					32,000	62										

IIT RESEARCH INSTITUTE

Table 7 (Cont.)

Date	Flight Route		Time of Day	Flight Time, min	Altitude, ft	Time at Altitude, min	Time Onset Detected, min							
	Origin	Destination					>1 PMH	>1 PMH	>1 PMH	>10 PMH	>15 PMH	>20 PMH	>25 PMH	>30 PMH
3/30/63	Keflavik	L. A.	D	557	35,000 36,000	285 210	240 210	195 210	195 210	15 135	15 90	15 25		
4/3/63	Chicago	London	N	470	33,000 37,000	51 369	30 369	139	240	160	85	15		
4/8/63	London	Seattle	D	569	32,000 35,000 40,000	43 247 225	35 247 225	35 120 225	25 95 225	25 80 225	10 55 190	20 150	65	15
4/19/63	Chicago	London	N	435	33,000 37,000	23 345	15 345	295	270	225	150	80		
4/22/63	London	Seattle	D	580	32,000 35,000 39,000 40,000	33 175 136 181	15 175 136 181	15 175 136 181	175 136 181	155 136 165	115 136 165	121 110	31 45	30
4/25/63	Chicago	London	N	450	33,000 35,000	166 232	166 232	140 145	125 115	55 95	5 80	40	10	
4/28/63	London	L. A.	D	662	32,000 35,000 36,000 39,000	35 238 147 187	10 238 147 187	10 238 110 187	10 238 110 187	10 238 110 135	165 165 85	95 20 25		
5/5/63	Chicago	London	D/N	411	33,000 34,000	343 18	290 18	120	120	50	20			
5/9/63	London	Seattle	D	563	35,000 36,000	193 299	193 292	193 292	193 292	193 292	163 227	163 147	23 47	
5/15/63	Seattle	Anchorage	N/D	177	39,000	121	121	121	121	121	106	95	80	
5/16/63	Anchorage	Chicago	D/N	328	29,000 33,000 34,000 37,000	34 71 77 105	34 71 77 105	25 71 77 105	25 71 77 105	25 71 77 60	71 77 45	65 77 40	35 40	30
5/17/63	Chicago	London	D/N	420	33,000 30,000	347 23	300 23	90	90	45				
5/23/63	London	Keflavik	D	146	28,000	109								
5/23/63	Keflavik	Seattle	D	435	35,000 36,000	285 95	285 95	265	225	155	130			
5/27/63	Chicago	London	D/N/D	446	33,000 30,000	352 28	352 28	50	35	5				

IIT RESEARCH INSTITUTE

Table 7 (Cont..)

Date	Flight Route		Time of Day	Flight Time, min	Altitude, ft	Time at Altitude, min	Time Since Detected, min									
	Origin	Destination					< 1 PMH	> 1 PMH	< 2.0 PMH	> 2.0 PMH	< 3.0 PMH	> 3.0 PMH	< 4.0 PMH	> 4.0 PMH	< 5.0 PMH	> 5.0 PMH
5/30/63	London	Chicago	D	515	32,000 35,000	23 439	23 325	115	95	40						
6/6/63	Chicago	London	D/W/D	461	32,000 37,000	95 297	60 297	297	252	237	237	122				
6/9/63	London	L. A.	D	641	28,000 35,000 36,000 37,000 40,000	16 210 160 20 190	60 115 15 190	60 115 190	60 115 190	30 95 175	75 75					
6/14/63	Chicago	London	D/W/D	447	29,000 37,000 39,000	32 136 224	60 160	60 160	60 160	30						
6/18/63	London	L. A.	D	690	32,000 33,000 35,000 36,000 39,000 40,000	24 115 136 82 147 134	20 115 85 65 30 134	20 100 85 50 134	20 100 85 50 134	10 20 30 30 134	5					
6/25/63	Chicago	London	D/W/D	444	30,000 33,000	22 373	22 225	22 120	22 90	10 40						
6/26/63	London	Keflavik	D	158	31,000 32,000	54 55	35 55	5	5							
6/28/63	Keflavik	L. A.	D	533	35,000 36,000 39,000	106 187 175	55 187 175	55 187 175	55 162 175	40 147 150	40 132 60					
7/8/63	Boston	London	W/D	365	31,000 35,000	65 228	5 45	5	5							
7/8/63	Chicago	London	D/W/D	445	29,000 30,000 33,000 34,000 35,000	11 1 129 32 210	13 13 129 32 210	13 65 140	13 65 125	13 55 75	50 32 60	50 20				
7/20/63	Anchorage	Seattle	W	170	37,000	124	124	124	124	124	65					
7/30/63	Chicago	Anchorage	D	350	35,000 36,000 39,000	51 143 105	120 105	120 105	120 105	120 75	55 40	55 10				

IIT RESEARCH INSTITUTE

Table 7 (Cont.)

Date	Flight Route		Time of Day	Flight Time, min.	Altitude, ft.	Time at Altitude, min.	Time Gases Detected, min.					
	Origin	Destination					>1 PMH	>2 PMH	>3 PMH	>4 PMH	>5 PMH	>10 PMH
8/1/63	Anchorage	Seattle	N	175	37,000	139	15	15	5			
8/16/63	Chicago	London	D/W/D	419	33,000	370	330	155	150	30		
8/21/63	London	Keflavik	D	151	32,000	111	5					
8/21/63	Keflavik	L. A.	D	526	31,000	81	15		15	5		
					35,000	169	90	25	15	5		
					36,000	219	140	125	95	10		

IIT RESEARCH INSTITUTE

Table 8
CHRONOLOGICAL SUMMARY OF OCSE DETECTION DURING FLIGHTS OVER POLAR LATITUDES

Date	Flight Route		Time of Day	Flight Time, min	Altitude, ft	Time at Altitude, min	Time OCSE Detected, min							
	Origin	Destination					>1 PPM	>3 PPM	>5 PPM	>10 PPM	>15 PPM	>20 PPM	>25 PPM	>30 PPM
7/20/63	Copenhagen	Anchorage	D	501	29,000 32,000 36,000 37,000	183 103 129 37	125 103 129 37	115 103 129 37	105 103 129 37	75 103 129 37	75 60 129 37	50 45 129 37		
7/31/63	Anchorage	Copenhagen	D	505	23,000 29,000 33,000 34,000 36,000	91 130 39 126 73	60 90 39 126 73	50 50 35 80 65	50 45 30 70 40	30 15 20 25 15				
8/3/63	Copenhagen	Anchorage	D	500	29,000 32,000 36,000 37,000	68 172 60 149	68 172 60 149	15 60 60 149	5 30 60 149	15 15 60 149	60 130	55 30		

IIT RESEARCH INSTITUTE

Table 9

CHRONOLOGICAL SUMMARY OF GEOME DETECTION DATA OBTAINED ON CONTRACT NO. FA 2668
(December 19, 1961, through April 30, 1962)

Date	Flight Route		Time of Day	Flight Time, min.	Altitude, ft.	Time at Altitude, min.	Time Geome Detected, min.				
	Origin	Destination					>1 PPM	>3 PPM	>5 PPM	>10 PPM	>20 PPM
2/6/62	S. F.	L. A.	M	75	27,000	10					
2/6/62	L. A.	S. F.	M	75	27,000	10					
2/7/62	S. F.	Chicago	D	225	28,000	30	20				
					33,000	45	45				
					37,000	120	100	70	20	20	5
2/16/62	Honolulu	L. A.	D/W	285	29,000	240					
2/17/62	L. A.	New York	M	270	29,000	105	60				
					31,000	135					
2/17/62	New York	Chicago	D	120	28,000	100					
2/25/62	Chicago	L. A.	M	240	28,000	180					
					31,000	30					
2/25/62	L. A.	Honolulu	M	120	28,000	140					
					31,000	160					
2/26/62	Honolulu	L. A.	D/M	260	37,000	210	45	25	10		
2/27/62	L. A.	Cleveland (New York)	M/D	350	33,000	120	120	75	75		
					37,000	180	180				
2/27/62	Cleveland	New York	D	65	27,000	45					
2/27/62	New York	S. F.	D	300	31,000	150	45			45	
					37,000	45	45	45	120	120	
2/27/62	S. F.	Seattle	M	100	35,000	55	45	45			
					37,000	145	105				
2/28/62	Seattle	Chicago	D	195	33,000	95	60	30	10		
3/5/62	Chicago	Phila.	D	120	35,000	95	55	45	45		
3/5/62	Phila.	Chicago	M	120	33,000	95	120	15			
3/18/62	Chicago	L. A.	M	240	28,000	200	195	195	110	75	
3/19/62	L. A.	New York	M	255	37,000	195	250				
3/19/62	New York	S. F.	D	325	28,000	275					
3/19/62	S. F.	Honolulu	D	295	37,000	255					

IIT RESEARCH INSTITUTE

Table 9 (Cont.)

Date	Origin	Destination	Time of Day	Flight Time, min	Altitude, ft	Time at Altitude, min	Time Ozone Detected, min				
							>1 PPM	>3 PPM	>5 PPM	>10 PPM	>15 PPM >20 PPM
3/20/62	Honolulu	S. F.	M	275	29,000	225	160	40	30		
3/20/62	S. F.	Chicago	D	210	29,000 33,000	85 110	5	50	50	15	
3/20/62	Chicago	Phila.	D	135	37,000	75	35	10			
3/20/62	Phila.	Chicago	M	145	31,000	75					
3/27/62	Chicago	L. A.	N	220	31,000	175					
3/27/62	L. A.	Baltimore	D	255	29,000 37,000	120 95		20	20	20	
3/27/62	Baltimore	L. A.	D/M	300	35,000 39,000	170 85	165 85	45 85	20 55	5	
3/28/62	L. A.	Chicago	M	200	37,000	165	165	60	50		
4/12/62	Chicago	Baltimore	D	90	31,000	35	35				
4/12/62	Baltimore	Chicago	D/M	120	28,000	60	55	5	5		
4/12/62	Chicago	Denver	N	115	33,000	75	75	30	30	30	
4/13/62	Denver	Chicago	D	110	29,000	65	65				
4/13/62	Chicago	Phila.	D	90	27,000 33,000	15 30	10 30	10 30	30	30	10
4/13/62	Phila.	Chicago	D/N	95	31,000	55	55	55	55	55	
4/15/62	Chicago	Baltimore	N	80	27,000	45	45	5			
4/16/62	Baltimore	Chicago	D	90	28,000	55	55	15	5		